

# Spin Physics in PIS

FRANCESCA GIORDANO 2011 RHIC & AGS ANNUAL USERS' MEETING











# Deep-inelastic scattering






longitudinal momentum structure




# Deep-inelastic scattering


FRANCESCA GIORDANO

2011 RHIC & AGS ANNUAL USERS' MEETING






longitudinal momentum structure




longitudinal spin structure




Deep-inelastic scattering





longitudinal momentum structure




longitudinal spin structure




only 30% of nucleon spin is carried by valence quarks




# Deep-inelastic scattering



























4	3			quark		
6			U	L		Т
n u	U	$f_1$	0		1	
c	L			g <sub>1</sub> 👶 -	8	
e o n	т				h <sub>1</sub> -(-	·






nucleon spin














4	8		quark		
6		U	L	Т	
n u	U	$f_1$ $\odot$			
c	L		g <sub>1</sub> 👶 - 🚷		
e o n	Т			h <sub>1</sub>	






nucleon spin

















A	3		quark	
6		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> 😲 - 🚷	
e o n	Т			h <sub>1</sub>





nucleon spin
















but integrated over final states!





A	£		quark	
O CO		U	L	Т
n u	U	$f_1$ $\odot$		
C	L		g <sub>1</sub> 👶 - 🚷	
e o n	Т			h <sub>1</sub>






nucleon spin













but integrated over final states!





A	3		quark	
<b>O</b>		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> 😲 - 🚷	
e o n	Т			h <sub>1</sub>





nucleon spin
















but integrated over final states!





4			quark	
6		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> • - •	
e o n	Т			h <sub>1</sub>

Transversity df

transverse longitudinal






nucleon spin













but integrated over final states!






A	3		quark	
O C		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> 😲 - 🚷	
e o n	т			h <sub>1</sub>

Transversity df inclusive 718

transverse longitudinal





nucleon spin
















but integrated over final states!





A	- Kin		quark	
6		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> 😲 - 🛞	
e o n	Т			h <sub>1</sub>

Transversity df inclusive 71S

transverse longitudinal






nucleon spin













but integrated over final states!






A	S.		quark	
6	<b>3</b>	U	L	Т
n u	U	$f_1$ $\odot$		
C	L		g <sub>1</sub> 😲 - 🛞	
e o n	Т			h <sub>1</sub>

Transversity df inclusive 71S

transverse longitudinal






nucleon spin













but integrated over final states!





A	S.		quark	
6	<b>3</b>	U	L	Т
n u	U	$f_1$ $\odot$		
C	L		g <sub>1</sub> 😲 - 🛞	
e o n	Т			h <sub>1</sub>

Transversity df inclusive 71S

 $h_1 \otimes X$ 

chiral odd chiral odd

transverse longitudinal






nucleon spin













but integrated over final states!





4			quark	
6		U	L	Т
n u	U	$f_1$ $\odot$		
c	L		g <sub>1</sub> 👶 - 🚷	
e o n	Т			h <sub>1</sub>

chiral odd

Transversity df




 $h_1 \otimes X$ 

chiral odd chiral odd

chiral even

transverse longitudinal






nucleon spin













 $A_{UT} \propto h_1 \otimes H_1^{\perp}$ 






$$A_{UT} \propto h_1 \otimes H_1^{\perp}$$
 Collins ff






 $A_{UT} \propto h_1 \otimes H_1^{\perp}$  Collins ff

#### 2-hadron production




 $A_{UT} \propto h_1 \times H_1^{\triangleleft}$ 






 $A_{UT} \propto h_1 \otimes H_1^{\perp}$  Collins ff

#### 2-hadron production




 $A_{UT} \propto h_1 imes H_1^{ riangleq}$  Interference ff






#### $A_{UT} \propto h_1 \otimes H_1^{\perp}$





# Collins mechanism: left-right asymmetry from correlations between quark transverse spin and outgoing hadron direction





































































#### Gaussian ansatz













#### Gaussian ansatz



















(different sign convention)



















#### 1-hadron production FRANCESCA GIORDANO






#### Sea role?

(different sign convention)








 $A_{UT} \propto h_1 \otimes H_1^{\perp}$  Collins ff


#### 2-hadron production



 $A_{UT} \propto h_1 imes H_1^{ riangleq}$  Interference ff

# Transversity in PIS







#### $A_{UT} \propto h_1 \times H_1^{\triangleleft}$


Interference Fragmentation:
left-right asymmetry from correlations between quark
transverse spin and relative orbital angular momentum of the
hadron pair
































FRANCESCA GIORDANO

2011 RHIC & AGS ANNUAL USERS' MEETING








$$A_{UT} \propto$$


$$A_{UT} \propto h_1 \times H_1^{\triangleleft}$$























#### Direct product! No assumption needed!













#### Direct product! No assumption needed!



in progress...





		quark				
		U		L	Т	
n u	U	$f_1$	0			
C	L			g <sub>1</sub> 👶 - 🚷		
e o n	т				h <sub>1</sub>	

transverse longitudinal





nucleon spin






parton spin







		quark				
		U	L	Т		
n u c l e o n	U	$f_1$ $\odot$		$h_1^{\perp}$ $\bigcirc$ - $\bigcirc$		
	L		g <sub>1</sub> 👶 - 🚷	$h_{1L}^{\perp}$ $\bigcirc$ - $\bigcirc$		
	Т	$f_{1T}^{\perp}$	$g_{1T}^{\perp}$ $\bullet$ $\bullet$ $\bullet$	$h_1$ $h_{1T}$ $h_{1T}$		

transverse longitudinal





nucleon spin







parton spin












transversity

#### transverse longitudinal





nucleon spin






parton spin


momentum











transverse longitudinal






nucleon spin






parton spin












$$\left\{ F_{\text{UU,T}} + \epsilon F_{\text{UU,L}} + \sqrt{2\epsilon (1+\epsilon)} \cos(\phi) F_{\text{UU}}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{\text{UU}}^{\cos(2\phi)} \right]$$

+ 
$$\lambda_l \left[ \sqrt{2\epsilon (1 - \epsilon)} \sin (\phi) F_{LU}^{\sin (\phi)} \right]$$

+ 
$$S_L$$
  $\left[\sqrt{2\epsilon (1 + \epsilon)} \sin (\phi) F_{UL}^{\sin (\phi)} + \epsilon \sin (2\phi) F_{UL}^{\sin (2\phi)}\right]$ 



worm-gears

Boer-Mulders  $\left[\sqrt{1-\epsilon^2}F_{\rm LL} + \sqrt{2\epsilon(1-\epsilon)}\cos(\phi)F_{\rm LL}^{\cos(\phi)}\right]$ 


+ 
$$S_T$$
  $\left[\sin(\phi - \phi_S)\left(F_{\text{UT},\text{T}}^{\sin(\phi - \phi_S)} + \epsilon F_{\text{UT},\text{L}}^{\sin(\phi - \phi_S)}\right)\right]$   
+  $\epsilon \sin(\phi + \phi_S)F_{\text{UT}}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S)F_{\text{UT}}^{\sin(3\phi - \phi_S)}$   
+  $\sqrt{2\epsilon(1 + \epsilon)}\sin(\phi_S)F_{\text{UT}}^{\sin(\phi_S)}$   
+  $\sqrt{2\epsilon(1 + \epsilon)}\sin(2\phi - \phi_S)F_{\text{UT}}^{\sin(2\phi - \phi_S)}$ 

+ 
$$S_T \lambda_l \left[ \sqrt{1 - \epsilon^2 \cos (\phi - \phi_S)} F_{LT}^{\cos (\phi - \phi_S)} + \sqrt{2\epsilon (1 - \epsilon)} \cos (\phi_S) F_{LT}^{\cos (\phi_S)} + \sqrt{2\epsilon (1 - \epsilon)} \cos (2\phi - \phi_S) F_{LT}^{\cos (2\phi - \phi_S)} \right] \right\}$$

transversity

pretzelosity

transverse longitudinal





nucleon spin





parton spin




n

Sivers






the parton transverse spin






the proton transverse spin




the parton transverse spin






the proton transverse spin




Both are Naively T-odd

the parton transverse spin






#### the proton transverse spin




the parton transverse spin



**Both are Naively T-odd** 




**Final State Interaction** 






#### the proton transverse spin




the parton transverse spin



Both are Naively T-odd




#### **Final State Interaction**






the proton transverse spin




the parton transverse spin




Both are Naively T-odd





**Final State Interaction** 






















#### Sivers asymmetries









Boer-Mulders asymmetries













Boer-Mulders asymmetries






unpolarized target & beam:








Boer-Mulders asymmetries








unpolarized target & beam:


spin effects also in unpolarized reactions!





Boer-Mulders asymmetries

X






$$\left\{ F_{\text{UU,T}} + \epsilon F_{\text{UU,L}} + \sqrt{2\epsilon (1+\epsilon)} \cos(\phi) F_{\text{UU}}^{\cos(\phi)} + \epsilon \cos(2\phi) F_{\text{UU}}^{\cos(2\phi)} \right]$$

+ 
$$\lambda_l \left[ \sqrt{2\epsilon (1 - \epsilon)} \sin (\phi) F_{LU}^{\sin (\phi)} \right]$$

+ 
$$S_L$$
  $\left[\sqrt{2\epsilon (1 + \epsilon)} \sin (\phi) F_{UL}^{\sin (\phi)} + \epsilon \sin (2\phi) F_{UL}^{\sin (2\phi)}\right]$ 



worm-gears

Boer-Mulders  $\left[\sqrt{1-\epsilon^2}F_{\rm LL} + \sqrt{2\epsilon(1-\epsilon)}\cos(\phi)F_{\rm LL}^{\cos(\phi)}\right]$ 


+ 
$$S_T$$
  $\left[\sin(\phi - \phi_S)\left(F_{\text{UT},\text{T}}^{\sin(\phi - \phi_S)} + \epsilon F_{\text{UT},\text{L}}^{\sin(\phi - \phi_S)}\right)\right]$   
+  $\epsilon \sin(\phi + \phi_S)F_{\text{UT}}^{\sin(\phi + \phi_S)} + \epsilon \sin(3\phi - \phi_S)F_{\text{UT}}^{\sin(3\phi - \phi_S)}$   
+  $\sqrt{2\epsilon(1 + \epsilon)}\sin(\phi_S)F_{\text{UT}}^{\sin(\phi_S)}$   
+  $\sqrt{2\epsilon(1 + \epsilon)}\sin(2\phi - \phi_S)F_{\text{UT}}^{\sin(2\phi - \phi_S)}$ 

+ 
$$S_T \lambda_l \left[ \sqrt{1 - \epsilon^2 \cos (\phi - \phi_S)} F_{LT}^{\cos (\phi - \phi_S)} + \sqrt{2\epsilon (1 - \epsilon)} \cos (\phi_S) F_{LT}^{\cos (\phi_S)} + \sqrt{2\epsilon (1 - \epsilon)} \cos (2\phi - \phi_S) F_{LT}^{\cos (2\phi - \phi_S)} \right] \right\}$$

transversity

pretzelosity

transverse longitudinal





nucleon spin






parton spin




n

Sivers













Pretzelosity








- \* chiral-odd > need the Collins FF or similar
- \* results consistent with zero at HERMES and COMPASS
- \* non-zero from CLAS?



#### worm gear UL










\* results consistent with zero \* hint for non-zero results?



worm gear LI







\* results consistent with zero \* hint for non-zero results?



worm gear LI

- \* To date, DIS experiments have played a crucial role in describing the proton structure
- \* Different reactions give complementary information: DIS, hadron and e+e-reactions are all needed to gain a complete picture of inner nucleon structure





- \* To date, DIS experiments have played a crucial role in describing the proton structure
- \* Different reactions give complementary information: DIS, hadron and e+e- reactions are all needed to gain a complete picture of inner nucleon structure

#### Thank you!



Summary